AD	

Award Number: DAMD17-00-1-0380

TITLE: The Use of Venezuelan Equine Encephalitis Replicons

Encoding the Her-2/neu Tumor Associated Antigen for the

Prevention and Treatment of Breast Cancer

PRINCIPAL INVESTIGATOR: Brian R. Long

Roland M. Tisch, Ph.D.

CONTRACTING ORGANIZATION: University of North Carolina

at Chapel Hill

Chapel Hill, North Carolina 27599-1350

REPORT DATE: May 2001

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. ACENICY LISE ONLY (Leave blank) 2. REPORT DATE

3. REPORT TYPE AND DATES COVERED

1. AGENCY USE ONLY (Leave blank)		Annual Summary			
	May 2001		. FUNDING NU		
4. TITLE AND SUBTITLE The Use of Venezuelan Equ Encoding the Her-2/neu Tu Prevention and Treatment	mor Associated Antigen fo	ns	DAMD17-00-		
6. AUTHOR(S)					
Brian R. Long					
Roland M. Tisch, Ph.D.					
7. PERFORMING ORGANIZATION NA University of North Carolina at C		8	B. PERFORMING REPORT NUM	GORGANIZATION MBER	
Chapel Hill, North Carolina 2759					
E-Mail: blong@med.unc.edu					
9. SPONSORING / MONITORING AG	GENCY NAME(S) AND ADDRESS(ES	3)	10. SPONSORING / MONITORING		
U.S. Army Medical Research and	Materiel Command		AGENCY RE	EPORT NUMBER	
Fort Detrick, Maryland 21702-50					
, ,					
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION / AVAILABILITY	STATEMENT	limitod		12b. DISTRIBUTION CODE	
Approved for Public Re.	lease; Distribution Un.	limitea	,		
13. ABSTRACT (Maximum 200 Wor	rds)				
14. SUBJECT TERMS				15. NUMBER OF PAGES	
Breast Cancer				7 16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFIC OF ABSTRACT	ATION	20. LIMITATION OF ABSTRACT	
Unclassified	Unclassified	Unclassifie	ed	Unlimited	

Table of Contents

•	Overview	Pg 1
•	Key Accomplishments	Pg 2
•	Reportable Outcomes	Pg 2
•	Figures	Pg 3
•	Summary	Pg 4
•	References	Pg 4

Overview:

During the past year, we have been focusing on the objectives outlined for Specific Aim 1 in our approved statement of work, and have completed an initial experiment related to Specific Aim 2. Testing and packaging of VEE replicon constructs encoding Her-2/neu, IL-12 and IL-18 have been completed. Additionally, a replicon construct encoding murine granulocyte-macrophage colony stimulating factor (mGM-CSF), has been established and tested and a truncated Her-2/neu gene encoding the extracellular and transmembrane domain of the protein is currently being assessed. We are hopeful the reduced size of theHer-2/neu gene will lead to increased levels of RNA and protein expression.

The rat Her-2/neu gene encodes for an 185kDa transmembrane protein. In order that we may have protein to work with experimentally, we have cloned overlapping fragments of the gene into a bacterial expression plasmid. Currently, one protein generated from the bacterial expression system, encompassing parts of the extracellular and transmembrane domains, has been used to demonstrate CD4⁺ T helper (Th) cell proliferation (Figure 1). For this experiment, mice were immunized intraperitoneally with 100µg of the protein fragment emulsified in incomplete Freund's adjuvant (IFA). Splenocytes were then recovered and assessed for T cell proliferation by ³H-thymidine uptake. Both rat HER-2/neu and A2-K^b transgenic mice showed significant proliferative responses. A second fragment, further spanning the extracellular domain, is in the process of being tested.

We have begun to characterize the CD8⁺ T cell response induced by replicons encoding Her-2/neu in both wild type FVB, and in FVB mice transgenic for the rat HER-2/neu gene. We have repeatedly generated significant CTL responses in wild type FVB mice, but CTL reactivity induced in Her-2/neu transgenic mice has to date been limited (Figure 2). The work of Jaffee et al. has demonstrated that Neu-specific T cell responses can be generated in Her-2/neu transgenic mice immunized with recombinant vaccinia virus expressing Neu. Consequently, we are hopeful that the appropriate combination of replicons encoding Her-2/neu as well as key inflammatory cytokines will overcome this element of immune tolerance seen in the transgenic mice.

One experiment has been completed in which both wild type FVB and Her-2/neu transgenic mice were immunized with replicons encoding Her-2/neu, and subsequently challenged with the F-H2N1 tumor cell line (expressing Her-2/neu). Both immunized and unimmunized Her-2/neu transgenic mice exhibited significant tumor growth whereas both groups of wild type mice rejected the tumor. While there was no significant difference in tumor weight from immunized and unimmunized Her-2/neu transgenic mice, decreased vascularity was noted in tumors recovered from the VEE Her-2/neu immunized mice. While we have no evidence that this decreased vascularity may be attributable to the vaccination, we remain optimistic that the correct combination of antigen and cytokine encoding replicons will be effective in inhibiting tumor growth.

Many of the tools are now in place that will allow us to test various antigen and cytokine combinations for optimal CD4⁺ Th and CD8⁺ CTL reactivity. This will further allow us to establish an effective approach of immunotherapy targeting Her-2/neu for the treatment and prevention of breast adenocarcinoma in transgenic mice.

Key Accomplishments:

• Completed in vitro testing and packaging of the following VEE replicons:

Full length Her-2/neu IL-12 IL-18 mGM-CSF IL-2

- Currently testing truncated Her-2/neu replicon encoding the extracellular and transmembrane domains.
- Cloned fragments of the Her-2/neu gene into a bacterial expression system and generated recombinant protein.
- Demonstrated CD4⁺ Th cell reactivity to one Her-2/neu protein fragment, and are in the process of testing a second fragment.
- Repeatedly generated CD8⁺ CTL reactivity in wild type FVB mice.
- Completed one tumor challenge experiment in VEE Her-2/neu immunized mice.

Reportable Outcomes:

• None.

Figures and Tables:




Figure 1. Proliferation in response to Her-2/neu fragment 4. Fragment 4 encompasses nucleotides 1856 – 2585 spanning part of the extracellular domain, the entire transmembrane domain and a small section of the intracellular domain. Her-2/neu transgenic and A2-K^b transgenic mice were twice immunized intraperitoneally with 100 μg fragment 4 emulsified in IFA. Splenocytes were harvested and incubated with the indicated concentration of recombinant protein and the levels of ³H-thymidine incorporation were measured. Data represents groups of three mice.

5 Hour 51 Cr Release Assay




Figure 2. Standard 5 hr chromium release assay for FVB and FVB/Neu^{+/+} mice immunized with VEE replicons encoding Her-2/neu (3.0x10⁵ IU) and replicons encoding mGM-CSF (3.0x10⁴ IU). Mice were immunized in the footpad 3X at 7 day intervals and sacrificed 3 – 5 days following the final injection. Spleen and popliteal lymph node cell suspensions were stimulated *in vitro* with psorelen inactivated NIH-3T3 cells (H-2^q) permanently transfected with Her-2/neu and B7-1 encoding constructs. Stimulated lymphocytes were incubated with 51 Cr labeled targets in the indicated ratios for 5 hours. 3T3 = non-transfected NIH-3T3 cells. Neu+B7 = NIH-3T3 cells transfected with Her-2/neu and B7.1. Data represents groups of 2 mice.

Summary:

Much of the work described for Specific Aim 1 has now been completed. We have finished the *in vitro* testing and packaging of the replicon constructs described, and have begun work on additional constructs. Progress has been made characterizing the CD8⁺ T cell response in both wild type and Her-2/neu transgenic mice with generation of consistent CTL activity in the wild type mice. Additionally, we have cloned overlapping fragments of the Her-2/neu gene into a bacterial expression system, and demonstrated a proliferative response to one of the recombinant proteins. These fragments will be used to evaluate CD4⁺ T helper responses in immunized mice. We believe an immunotherapeutic protocol consisting of the appropriate combination and antigen and cytokine encoding replicons will be useful for overcoming the tolerance associated with the Her-2/neu transgenic mice.

In our initial tumor challenge experiment with replicons encoding Her-2/neu alone, we failed to mediate rejection of transferred Neu expressing tumors in Her-2/neu transgenic mice. Again, we feel it will be a matter of determining the appropriate cytokine encoding constructs for use in our protocol.

References:

Reilly, R. Todd, Morris B.C. Gottlieb, Anne M. Ercolini, Jean-Pascal H. Machiels, Colin E. Kane, Francesca I. Okoye, William J. Muller, Katherine H. Dixon, and Elizabeth M. Jaffee. (2000). Her-2/neu Is a Tumor Rejection Target in Tolerized HER-2/neu Transgenic Mice. Cancer Research 60, 3569-3576